

# Return to School Plan-Home Concussion Management (Appendix E)

The Initial Rest to Step 2 of the Return to School Plan is the Home Concussion Management Plan

The Home Concussion Management Plan is the first two steps before a student, that has been diagnosed with a concussion, can start the Return to School Plan. The following steps will be followed if the student has received a diagnosis of concussion from doctor or nurse practitioner. Medical Concussion Assessment (Appendix D Part 2) would have had to be completed and signed off by a doctor or nurse practitioner. The parent/guardian/caregiver monitor and sign off on each step.

Each step must last a minimum of 24 hours.

#### **Initial Rest**

Focus: 24 48 hours of relative cognitive rest:

Activities permitted if tolerated by student:

short board/card g@en)e2[E)2 11.04 Tf@aml9.0000092 0 62;[ho)3(r)-3(t)6)-4(bo)3(ard)10(/)-4(card)10



# Step 1

# Focus:

 light cognitive (thinking/memory/ knowledge) activities; and gradually increase cognitive activity up to 30 minutes. Take frequent breaks.

Activities permitted if tolerated by student:

activities from previous stage; easy reading (for example, books, magazines, newspaper); limited tv; limited cellphone conversations; drawing/building blocks/puzzles; and some contact with friends.

Activities that are NOT permitted at this stage:

technology use (for example, computer, laptop, tablet, cell phone; and attendance at school or school type work.

### The student moves to Step 2 when:

the student tolerates 30-minutes of light cognitive activity (for example a student should be able to complete 3-4 of the permitted activities in Step 1) and has not exhibited or reported a return of symptoms, new symptoms, or worsening symptoms; and

the student has completed a minimum of 24-hours at Step 1.

#### However:

the student must return to the previous step for a minimum of 24-hours when they have exhibited or reported a return of symptoms, or new symptoms.

At any time, the student must return to a medical doctor or nurse practitioner when they have exhibited or reported a worsening of symptoms